arXiv:hep-th/9812238v1 28 Dec 1998

SMI-25-98

Note on the Massive Rarita-Schwinger Field in the AdS/CFT Correspondence

A. S. Koshelev[§] and O. A. Rytchkov[⋆],

[§] *Department of Theoretical Physics, Moscow State University,*
Moscow, Russia, 119899
kas@depni.npi.msu.su

[⋆] *Steklov Mathematical Institute,*
Gubkin str.8, Moscow, Russia, 117966
rychkov@orc.ru

Abstract

We consider a massive Rarita-Schwinger field on the Anti-de Sitter space and solve the corresponding equations of motion. We show that appropriate boundary terms calculated on-shell give two-point correlation functions for spin-3/2 fields of the conformal field theory on the boundary. The relation between Rarita-Schwinger field masses and conformal dimensions of corresponding operators is established.

The correspondence (duality) between supergravity on $d+1$-dimensional Anti-de Sitter space and a certain superconformal field theory on its d-dimensional boundary, proposed in [1] and elaborated in [2, 3] has already undergone a large number of examinations. The most important example of this correspondence is provided by IIB supergravity on $AdS_5 \times S^5$ background which is dual to $\mathcal{N} = 4$, $D = 4$ SYM, appearing on the AdS_5 boundary. Another pair of dual theories involve eleven-dimensional supergravity compactified on $AdS_7 \times S^4$ and (2,0) superconformal field theory on the AdS_7 boundary.

According to this correspondence all physical exitations of the bulk supergravity can be put in one-to-one correspondence with the composite operators of the conformal theory on the boundary and the generating functional for these operators is equal to the exponentiated classical supergravity action calculated on-shell with the given boundary conditions specified by the function ϕ_0

$$\left\langle \exp\left(\int_{\partial AdS_{d+1}} d^d x \mathcal{O}\phi_0\right)\right\rangle_{CFT} = \exp(-I(\phi)). \tag{1}$$

Thus in order to find the correlation functions of operators in conformal field theory one has to solve the classical Dirichlet problem for the corresponding supergravity field on the AdS space. It gives the direct way to check the correspondence: one should compare the correlation functions expected in conformal field theories and those obtained via supergravity. In this way two-point functions of local operators corresponding to the scalar fields [3, 4, 5, 6, 7], vector fields [3, 6, 8, 9], spinor fields [10, 8], metric tensor fields [11, 12, 13], antisymmetric form fields [14, 15, 16] and massless Rarita-Schwinger fields [17, 18] were investigated and the agreement with the conformal field results was established. Also various three-point correlation functions for these fields were examined [4, 5, 6, 8, 19, 20]. In all cases the results are in agreement with those expected in the conformal field theory. In this note we consider the massive Rarita-Schwinger field and study its two-point correlation function.

The general action for the massive Rarita-Schwinger field Ψ_μ in a $d+1$-dimensional curved space is [21]

$$S = \int d^{d+1}x\sqrt{g}(\bar{\Psi}_\mu \Gamma^{\mu\nu\rho} D_\nu \Psi_\rho - m_1 \bar{\Psi}_\mu \Psi^\mu - m_2 \bar{\Psi}_\mu \Gamma^{\mu\nu}\Psi_\nu). \tag{2}$$

Here D_μ is a covariant derivative, $\Gamma^{\mu\nu\rho} = \Gamma^{[\mu}\Gamma^{\nu}\Gamma^{\rho]}$, $\Gamma^{\mu\nu} = \Gamma^{[\mu}\Gamma^{\nu]}$, where Γ_μ are curved space gamma matrices connected with the flat space gamma matrices γ_a via vielbein e^a_μ: $\Gamma_\mu = e^a_\mu \gamma_a$, μ, $a = 0, 1, .., d$. The flat space gamma matrices are subjects of the anticommutation relations $\{\gamma_a, \gamma_b\} = 2\delta_{ab}$. We suppose that the background is $d+1$-dimensional Euclidean Anti-de Sitter space with the coordinates $x = (x_0, \vec{x}) = (x_0, x_i)$, where $x_0 > 0$, $i = 1, .., d$. The hypersurface $x_0 = 0$ will be considered as the AdS boundary. The corresponding AdS metric has the form

$$ds^2 = \frac{1}{x_0^2}(dx_0^2 + dx_i dx^i) \tag{3}$$

and we can specify the vielbein as

$$e^a_\mu = \frac{1}{x_0}\delta^a_\mu. \tag{4}$$

Below we will investigate the action (2) with rather general mass parameters m_1 and m_2. However the most interesting results are related with the Rarita-Schwinger field on AdS_5 appearing in the compactification of IIB supergravity on $AdS_5 \times S^5$. The corresponding mass spectrum was found in [22] and consists of three sets of states:

$m_1 = -(3/2 + k), \quad m_2 = 0, \quad k = 0, 1, 2, ..$
$m_1 = 7/2 + k, \quad m_2 = 0, \quad k = 0, 1, 2, ..$
$m_1 = 0, \quad m_2 = -3/2.$

The action (2) leads to the following equation of motion

$$\Gamma^{\mu\nu\rho} D_\nu \Psi_\rho - m_1 \Psi^\mu - m_2 \Gamma^{\mu\nu} \Psi_\nu = 0, \tag{5}$$

which can be rewritten in the equivalent form

$$\Gamma^\nu (D_\nu \Psi_\rho - D_\rho \Psi_\nu) - m_- \Psi_\rho + \frac{m_+}{d-1} \Gamma_\rho \Gamma^\nu \Psi_\nu = 0, \tag{6}$$

where $m_\pm = m_1 \pm m_2$. For the future convenience we project the Rarita-Schwinger field on the vielbein introducing

$$\psi_a = e^\mu_a \Psi_\mu. \tag{7}$$

Substitution of this field and vielbein (4) in Eq. (6) is straightforward and gives

$$\begin{gathered} x_0 \gamma^a \partial_a \psi_b - x_0 \partial_b (\gamma^a \psi_a) - \gamma_b \psi_0 - \frac{d}{2} \gamma_0 \psi_b - \frac{1}{2} \gamma_0 \gamma_b (\gamma^a \psi_a) \\ + \frac{3}{2} \delta_{0b} (\gamma^a \psi_a) - m_- \psi_b + \frac{m_+}{d-1} \gamma_b (\gamma^a \psi_a) = 0. \end{gathered} \tag{8}$$

Following Witten [3], in order to solve these equations we will find a solution which depends only on x_0 coordinate and has a singularity on the infinity and then apply the inversion (accompanied by the rotation of the vielbein), which moves the singularity to the boundary $x_0 = 0$. The fields depending only on x_0 coordinate are the subject of the following equations

$$x_0 \gamma^0 \partial_0 \psi_0 - (\frac{d}{2} + 1) \gamma_0 \psi_0 - m_- \psi_0 = x_0 \partial_0 \chi - \frac{m_+}{d-1} \gamma_0 \chi - \chi, \tag{9}$$

$$x_0 \gamma^0 \partial_0 \psi_i - \frac{d}{2} \gamma_0 \psi_i - m_- \psi_i = \frac{1}{2} \gamma_0 \gamma_i \chi - \frac{m_+}{d-1} \gamma_i \chi + \gamma_i \psi_0, \tag{10}$$

where we denote $\chi = \gamma^a \psi_a$. It is easy to see that these equations put the constraints on the fields

$$(d - 1 - 2m_- \gamma_0) \psi_0 = (d - 1 + 2m_2 \gamma_0) \chi. \tag{11}$$

Taking (11) into account we get the following solution for the equations (9), (10)

$$\psi_0 = x_0^{d/2+C} b_0^+ + x_0^{d/2-C} b_0^-, \tag{12}$$

$$\begin{gathered} \psi_i = x_0^{d/2+m_-} b_i^+ + x_0^{d/2-m_-} b_i^- - \\ - \frac{2m_1}{d(d-1-2m_2)} x_0^{d/2-C} \gamma_i b_0^- - \frac{2m_1}{d(d-1+2m_2)} x_0^{d/2+C} \gamma_i b_0^+, \end{gathered} \tag{13}$$

where

$$C = \frac{d(d-1)}{4m_1} + \frac{(m_1 - m_2)(m_1 + dm_2)}{m_1(d-1)} \tag{14}$$

and $b_0^{\pm}$, $b_i^{\pm}$ are constant spinors with a definite chirality ($\gamma_0 b_a^+ = b_a^+$, $\gamma_0 b_a^- = -b_a^-$). Also $b_0^{\pm}$, $b_i^{\pm}$ are the subject of the additional relations

$$\gamma^i b_i^+ = 0, \quad \gamma^i b_i^- = 0. \tag{15}$$

Applying O(d+1,1) transformation of the AdS_{d+1} isometry group we can map infinity to an arbitrary point on the boundary $x_0 = 0$. It introduces the dependence on x_i coordinates. The corresponding transformation is

$$x_\mu \longrightarrow \frac{x_\mu}{x_0^2 + \vec{x}^2}. \tag{16}$$

Note that an application of this transformation changes the vielbein (4), so in order to get a solution in an old local Lorentz frame we have to transform a vielbein as follows

$$e_\mu^a \longrightarrow L_b^a e_\mu^b, \quad \text{where} \quad L_b^a = \delta_b^a - \frac{2x^a x_b}{x_0^2 + \vec{x}^2}. \tag{17}$$

This inversion of a vielbein is supplemented by an appropriate transformation of the spinor field [24]

$$\psi_a \longrightarrow L_a^b \frac{x_0\gamma^0 + x_i\gamma^i}{\sqrt{x_0^2 + \vec{x}^2}} \psi_b. \tag{18}$$

Note that this transformation changes the sign of gamma-matricies so in order to recover the invariance of equations of motion we have to change all mass signs. The above procedure, applied to the solution (12), (13) gives us bulk-to-boundary propagator for the gravitino field. The contraction of this propagator with arbitrary functions yields

$$\begin{aligned}
\psi_0 = \int d^d y \Bigg(& \frac{(x_0\gamma^0 + (x-y)_i\gamma^i) x_0^{d/2 - C\gamma_0}}{(x_0^2 + (\vec{x}-\vec{y})^2)^{(d+1)/2 - C\gamma_0}} (1 - \frac{2x_0^2}{x_0^2 + (\vec{x}-\vec{y})^2}) b_0(\vec{y}) \\
& - \frac{2(x_0\gamma^0 + (x-y)_i\gamma^i) x_0^{d/2+1-m_-\gamma_0}}{(x_0^2 + (\vec{x}-\vec{y})^2)^{(d+1)/2+1-m_-\gamma_0}} (x-y)_i b_i(\vec{y}) \\
& - \frac{4m_1}{d(d-1+2m_2)} \frac{(x_0\gamma^0 + (x-y)_i\gamma^i) x_0^{d/2+1+C}}{(x_0^2 + (\vec{x}-\vec{y})^2)^{(d+1)/2+1+C}} (x-y)_i \gamma_i b_0^-(\vec{y}) \\
& - \frac{4m_1}{d(d-1-2m_2)} \frac{(x_0\gamma^0 + (x-y)_i\gamma^i) x_0^{d/2+1-C}}{(x_0^2 + (\vec{x}-\vec{y})^2)^{(d+1)/2+1-C}} (x-y)_i \gamma_i b_0^+(\vec{y}) \Bigg),
\end{aligned} \tag{19}$$

$$\begin{aligned}
\psi_i = \int d^d y \Bigg(& \frac{(x_0\gamma^0 + (x-y)_i\gamma^i) x_0^{d/2 - m_-\gamma_0}}{(x_0^2 + (\vec{x}-\vec{y})^2)^{(d+1)/2 - m_-\gamma_0}} (\delta_{ij} - \frac{2(x-y)_i(x-y)_j}{x_0^2 + (\vec{x}-\vec{y})^2}) b_j(\vec{y}) \\
& - \frac{2(x_0\gamma^0 + (x-y)_i\gamma^i) x_0^{d/2+1-C\gamma_0}}{(x_0^2 + (\vec{x}-\vec{y})^2)^{(d+1)/2+1-C\gamma_0}} (x-y)_i b_0(\vec{y})
\end{aligned}$$

$$+\frac{2m_1}{d(d-1+2m_2)}\frac{(x_0\gamma^0+(x-y)_i\gamma^i)x_0^{d/2+C}}{(x_0^2+(\vec{x}-\vec{y})^2)^{(d+1)/2+C}}(\delta_{ij}-\frac{2(x-y)_i(x-y)_j}{x_0^2+(\vec{x}-\vec{y})^2})\gamma_j b_0^-(\vec{y})$$

$$+\frac{2m_1}{d(d-1-2m_2)}\frac{(x_0\gamma^0+(x-y)_i\gamma^i)x_0^{d/2-C}}{(x_0^2+(\vec{x}-\vec{y})^2)^{(d+1)/2-C}}(\delta_{ij}-\frac{2(x-y)_i(x-y)_j}{x_0^2+(\vec{x}-\vec{y})^2})\gamma_j b_0^+(\vec{y})\Bigg). \qquad (20)$$

Eq. (19) and (20) give the general solution to the equations of motion (8). In the same way we construct the solution for the conjugated spinors $\bar{\psi}_a$, that has the same form except the change of all mass signs.

Now we have to impose the appropriate boundary conditions on the infinity and on the hypersurface $x_0 = 0$. Namely, in the solution (19), (20) we keep only the terms, which vanish as x_0 goes to infinity. Note that the asymptotic behavior of the fields essentially depends on the values of the constants d, m_1 and m_2. Moreover, different terms in (19), (20) have different limits as x_0 tends to zero. In order to be concrete in all formulae below we suppose m_1 and $m_1 - m_2$ to be positive (all other cases could be studied in the same way). We denote $\psi_a = \psi_a^{'}[b_i^{\pm}] + \psi_a^{''}[b_0^{\pm}]$, where we collect in $\psi_a^{'}$ all terms depending on b_i and in $\psi_a^{''}$ all terms depending on b_0. The fields $b_i^{\pm}(\vec{y})$ define the asymptotics of $x_0^{-d/2+m_-}\psi_a^{'}[b_i^{\pm}]$. As in the Dirac case [10] in order to obtain a square integrable asymptotics on the boundary we have to put $b_i^+(\vec{y}) = 0$. The asymptotic behavior of the terms $x_0^{-d/2+m_-}\psi_a^{''}[b_0^{\pm}]$ is determined by the relation between two constants m_- and C. In general, due to these relations the functions b_0 couldn't be fixed by the boundary data, since all the terms $x_0^{-d/2+m_-}\psi_a^{''}[b_0^{\pm}]$ vanish or go to infinity as x_0 approaches zero. Thus only b_i^- are determined by the boundary conditions. Note that in the particular case $b_0 = 0$ we have the following solution to the Dirichlet problem for the Rarita-Schwinger field on the AdS_{d+1} space

$$\psi_0(x_0,\vec{x}) = -\int d^dy\frac{2(-x_0+(x-y)_i\gamma^i)x_0^{d/2+1+m_-}}{(x_0^2+(\vec{x}-\vec{y})^2)^{(d+1)/2+1+m_-}}(x-y)_i b_i^-(\vec{y}), \qquad (21)$$

$$\psi_i(x_0,\vec{x}) = \int d^dy\frac{(-x_0+(x-y)_i\gamma^i)x_0^{d/2+m_-}}{(x_0^2+(\vec{x}-\vec{y})^2)^{(d+1)/2+m_-}}\left(\delta_{ij}-\frac{2(x-y)_i(x-y)_j}{x_0^2+(\vec{x}-\vec{y})^2}\right)b_j^-(\vec{y}). \qquad (22)$$

The general scheme of AdS/CFT correspondence [3] prescribes us to calculate the classical action on-shell with given boundary conditions. The action for the Rarita-Schwinger field (2) is linear so it vanishes on-shell. As in the Dirac case [10] it should be supplied by a special boundary term, which should be covariant and quadratic in the Rarita-Schwinger fields [17, 18]. We take such term in the form $I = I_1 + I_2$, where

$$I_1 = \mathrm{const}\cdot\lim_{\varepsilon\to 0}\int_{M_d^{\varepsilon}} d^dx\sqrt{g_{\varepsilon}}\bar{\Psi}_i(\vec{x})g_{\varepsilon}^{ij}\Psi_j(\vec{x}), \qquad (23)$$

$$I_2 = \mathrm{const}\cdot\lim_{\varepsilon\to 0}\int_{M_d^{\varepsilon}} d^dx\sqrt{g_{\varepsilon}}\bar{\Psi}_i(\vec{x})\Gamma^i\Gamma^j\Psi_j(\vec{x}). \qquad (24)$$

M_d^{ε} is the hypersurface $x_0 = \varepsilon$, $\sqrt{g_{\varepsilon}}$ is a volume element in the induced metric g_{ij} on this hypersurface. The second structure (24) doesn't contribute to the correlation function of operators coupled to b_i^- since on-shell we have $\gamma^a\psi_a \sim b_0$. In order to calculate I_1

we will substitute (20) and the corresponding solution for the conjugated spinor in (23) and rewrite it as a sum of the integrals with the similar structure. So the integral to be computed is

$$G(y_1, y_2, \epsilon) = \int d^d x \varepsilon^{-d} \left(\delta_{im} - 2\frac{(x-y_1)_i (x-y_1)_m}{\varepsilon^2 + (\vec{x}-\vec{y}_1)^2} \right) \times$$
$$\times \left(\frac{\varepsilon\gamma^0 + (x-y_1)_k \gamma^k}{(\varepsilon^2 + (\vec{x}-\vec{y}_1)^2)^{d/2+1/2+\sigma}} \varepsilon^{d/2+\sigma} \right) \left(\frac{\varepsilon\gamma^0 + (x-y_2)_j \gamma^j}{(\varepsilon^2 + (\vec{x}-\vec{y}_2)^2)^{d/2+1/2+\rho}} \varepsilon^{d/2+\rho} \right) \times$$
$$\times \left(\delta_{in} - 2\frac{(x-y_2)_i (x-y_2)_n}{\varepsilon^2 + (\vec{x}-\vec{y}_2)^2} \right). \tag{25}$$

In Appendix we argue that this integral is singular in the limit $\epsilon \longrightarrow 0$ if the difference between ρ and σ is not integer. Except very special cases of masses the difference between C and m_- is not integer, so the terms mixing b_0 and b_i give a singular contribution in I_1. Therefore if we perform the "minimal" renormalization [4], i.e. just drop out singular terms, we get that the generating functional doesn't depend on the terms containing simultaneously b_0 and b_i. So the correlation functions of operators coupled to b_i are determined by the term involving only b_i. In order to calculate the latter one we will also use the results of Appendix. In this case the hypergeometric function ${}_3F_2$ in (36) turns to ${}_2F_1$, so we have

$$\int \frac{\varepsilon^{2m_-+1}}{(\varepsilon^2 + (\vec{x}-\vec{y}_1)^2)^{(d+1)/2+m_-} (\varepsilon^2 + (\vec{x}-\vec{y}_2)^2)^{(d+1)/2+m_-}} d^d x$$
$$= \frac{\text{const}}{\varepsilon^{d+2m_-+1}} {}_2F_1 \left(\frac{d}{2} + 1 + 2m_-, \frac{d}{2} + \frac{1}{2} + m_-; \frac{d}{2} + m_- + 1; -\frac{|\vec{y}_1 - \vec{y}_2|^2}{4\varepsilon^2} \right)$$
$$= \frac{\text{const}}{(4\varepsilon^2 + (\vec{y}_1 - \vec{y}_2)^2)^{(d+1)/2+m_-}} {}_2F_1 \left(\frac{d}{2} + \frac{1}{2} + m_-, -m_-; \frac{d}{2} + m_- + 1; \frac{|\vec{y}_1 - \vec{y}_2|^2}{4\varepsilon^2 + |\vec{y}_1 - \vec{y}_2|^2} \right) \tag{26}$$

The result for the integral turns out to be

$$I_1 = \text{const} \cdot \int \ d^d y_1 d^d y_2 \bar{b}_i^+ (\vec{y}_1) \frac{(y_1 - y_2)_k \gamma^k}{|\vec{y}_1 - \vec{y}_2|^{d+1+2m_-}} (\delta_{ij} - \frac{2(y_1-y_2)_i (y_1-y_2)_j}{(\vec{y}_1 - \vec{y}_2)^2}) b_j^- (\vec{y}_2). \tag{27}$$

As it was noted in [4, 7] the integrals obtained after calculation the actions on-shell and taking the limit $\epsilon \longrightarrow 0$ are divergent and need the regularization. It was suggested to make renormalization or to consider the integral (27) as a value of the distribution

$$\frac{(y_1 - y_2)_k \gamma^k}{|\vec{y}_1 - \vec{y}_2|^{d+1+2m_-}} (\delta_{ij} - \frac{2(y_1-y_2)_i (y_1-y_2)_j}{(\vec{y}_1 - \vec{y}_2)^2}) \tag{28}$$

on a test function. The Fourier transform of this distribution is

$$F\left[\frac{y_k \gamma^k}{|\vec{y}|^{d+2m_-+1}} (\delta_{ij} - \frac{2 y_i y_j}{\vec{y}^2}) \right] = \text{const} \cdot p^{2m-1} p_k \gamma^k \left(\delta_{ij} + \frac{2}{(d+2m_- - 3)(d+2m_- - 1)} \frac{p_i p_j}{p^2} \right) \tag{29}$$

for $2m_- + 1 \neq 2k$, where $k = 0, 1, ..$ and $p = |\vec{p}|$. The overall factor in (29) is not important for us. Also we take in account that the test functions in (27) are the subject of the restriction $\gamma^i b_i = 0$. When $2m_- + 1$ is even, the Fourier transform includes a logarithmic term

$$F\left[\frac{y_k\gamma^k}{|\vec{y}|^{d+2m_-+1}}(\delta_{ij} - \frac{2y_iy_j}{\vec{y}^2})\right]$$

$$= \mathrm{const} \cdot p^{2m-1}\log\frac{p}{c}p_k\gamma^k\left(\delta_{ij} + \frac{2}{(d+2m_- - 3)(d+2m_- - 1)}\frac{p_ip_j}{p^2}\right). \qquad (30)$$

As it was noted in [7] the distributions of this type are not conformal invariant.

The general scheme prescribes to variate the unconstrained fields on the boundary, however, we have a restriction $\gamma_i b_i(\vec{y}) = 0$. It means that we should introduce a projection operator on this constraint, namely

$$b_i(\vec{y}) = (\delta_{ij} - \frac{1}{d}\gamma_i\gamma_j)\tilde{b}_j(\vec{y}) = \Pi_{ij}\tilde{b}_j(\vec{y}), \qquad (31)$$

where $\tilde{b}_j(\vec{y})$ is an unconstrained field. Taking it into account and using (27) we obtain

$$\left\langle\Sigma^+_{i\alpha}(\vec{y}_1)\Sigma^-_{j\beta}(\vec{y}_2)\right\rangle = \mathrm{const} \cdot \Pi_{ik}\frac{(y_1-y_2)_i(\gamma^i)_{\alpha\beta}}{|\vec{y}_1-\vec{y}_2|^{d+1+2m_-}}(\delta_{kl} - \frac{2(y_1-y_2)_k(y_1-y_2)_l}{(\vec{y}_1-\vec{y}_2)^2})\Pi_{lj}, \qquad (32)$$

$\Sigma_{i\alpha}(\vec{y})$ is a conformal field operator which corresponds to the Rarita-Schwinger field in supergravity. Note that the second projection operator in (32) could be eliminated by the trivial algebraic transformations, so only the first one is really necessary. The obtained correlation function coincides with the projected two-point correlation function expected in the conformal field theory for operators with spin-3/2 [26] and with the conformal dimension

$$\Delta = \frac{d}{2} + m_-. \qquad (33)$$

For the massless case $m_1 = 0$ these results coincide with those obtained in [17, 18].

In this note we have considered the action for the massive Rarita-Schwinger field in the Anti-de Sitter space and solve the corresponding equations of motion. According to the prescriptions of AdS/CFT correspondence we have demanded the appropriate boundary behavior of the field and using the special boundary terms we have obtained the two-point correlation function for the conformal field theory operators with spin-3/2. In our consideration we didn't specify the AdS dimension d and the parameters m_1 and m_2 of the Rarita-Schwinger field. However we implied that these parameters are subject of the special relations: we supposed that the difference between C and m_- is not integer. It is easy to see that for the Rarita-Schwinger field masses in IIB supergravity on $AdS_5 \times S^5$ [22] and in eleven-dimensional supergravity on $AdS_7 \times S^4$ [23] the above condition is satisfied.

ACKNOWLEDGEMENTS

We are grateful to I. Ya. Aref'eva for suggesting this problem and permanent interest to our work. Also we are grateful to G. E. Arutyunov for useful and stimulating discussions. A.K. and O.R. are supported in part by RFFI grant 96-01-00608. O.R. is supported in part by the INTAS Grant 96-0457 within the research program of the International Center for Fundamental Physics in Moscow.

Appendix

The purpose of this Appendix is to give an explicit calculation of the general integral (25). Performing multiplication, one can get

$$G(y_1,y_2,\epsilon)=\int d^dx\,\varepsilon^{\rho+\sigma}\frac{\varepsilon^2+(x-y_1)_k\gamma^k(x-y_2)_j\gamma^j-\varepsilon(y_1-y_2)_k\gamma^k\gamma_0}{(\varepsilon^2+(\vec{x}-\vec{y}_1)^2)^{d/2+1/2+\sigma}(\varepsilon^2+(\vec{x}-\vec{y}_2)^2)^{d/2+1/2+\rho}}\times$$
$$\times\left(\delta_{mn}+4\frac{(x-y_1)_i(x-y_2)^i(x-y_1)_m(x-y_2)_n}{(\varepsilon^2+(\vec{x}-\vec{y}_1)^2)(\varepsilon^2+(\vec{x}-\vec{y}_2)^2)}\right.$$
$$\left.-2\frac{(x-y_1)_m(x-y_1)_n}{\varepsilon^2+(\vec{x}-\vec{y}_1)^2}-2\frac{(x-y_2)_m(x-y_2)_n}{\varepsilon^2+(\vec{x}-\vec{y}_2)^2}\right). \tag{34}$$

We will consider (34) as a sum of different terms with the similar structure, it will unify our investigation. Also the succeeding analysis will be simplified if we rewrite each term applying the Fourier transformation. We use the following Fourier transformation formula:

$$2^\alpha\Gamma\left(\frac{d}{2}+\alpha\right)\frac{\varepsilon^\alpha}{(\varepsilon^2+(\vec{x}-\vec{y})^2)^{d/2+\alpha}}=\mathrm{const}\cdot\int d^dp\,e^{i\vec{p}(\vec{x}-\vec{y})}p^\alpha K_\alpha(\varepsilon p) \tag{35}$$

where K_α is a modified Bessel function with index α and $p=|\vec{p}|$. The contraction of two propagators in (34) turns into the product of their Fourier images. The remaining integral could be found explicitly in [27], so we get

$$\int e^{-i\vec{p}(\vec{y_1}-\vec{y_2})}p^{\mu+\nu}K_\mu(\varepsilon p)K_\nu(\varepsilon p)d^dp=$$
$$=\frac{\mathrm{const}}{\varepsilon^{d+\mu+\nu}}\frac{2^{d/2+\mu+\nu-2}\Gamma\left(d/2+\mu+\nu\right)\Gamma\left(d/2+\nu\right)\Gamma\left(d/2+\mu\right)}{\Gamma\left(d+\mu+\nu\right)}\times$$
$${}_3F_2\left(d/2+\mu+\nu,d/2+\nu,d/2+\mu;\frac{d+\mu+\nu}{2},\frac{d+\mu+\nu+1}{2};-\frac{|\vec{y}_1-\vec{y}_2|^2}{4\varepsilon^2}\right). \tag{36}$$

All factors like p_m appearing after differentiation of (35) could be obtained by applying the operator $i\frac{\partial}{\partial y_m}$ to (36). The circle of convergence for the hypergeometric function ${}_3F_2(z)$ is $|z|<1$. So, in order to find the limit $\varepsilon\to 0$ we have to apply the formula of analytic continuation, known in the theory of the hypergeometric functions

$${}_3F_2(a_1,a_2,a_3;b_1,b_2;z)=\frac{\Gamma(b_1)\Gamma(b_2)}{\Gamma(a_1)\Gamma(a_2)\Gamma(a_3)}\left[\Gamma(a_1)(-z)^{-a_1}\frac{\Gamma(a_2-a_1)\Gamma(a_3-a_1)}{\Gamma(b_1-a_1)\Gamma(b_2-a_1)}\times\right.$$
$${}_3F_2(1+a_1-b_1,1+a_1-b_2,a_1;1+a_1-a_2,1+a_1-a_3;1/z)$$

$$+\Gamma(a_2)(-z)^{-a_2}\frac{\Gamma(a_1-a_2)\Gamma(a_3-a_2)}{\Gamma(b_1-a_2)\Gamma(b_2-a_2)}\times$$

$$\times{}_3F_2(1+a_2-b_1,1+a_2-b_2,a_2;1+a_2-a_1,1+a_2-a_3;1/z)$$

$$+\Gamma(a_3)(-z)^{-a_3}\frac{\Gamma(a_1-a_3)\Gamma(a_2-a_3)}{\Gamma(b_1-a_3)\Gamma(b_2-a_3)}\times$$

$$\times\,{}_3F_2(1+a_3-b_1,1+a_3-b_2,a_3;1+a_3-a_1,1+a_3-a_2;1/z)\Big]\,. \tag{37}$$

Taking into account, that $F(0)=1$ we find three terms with the different behavior as ε tends to zero. In our concrete case of the integral (34) we have $\mu=\sigma+1/2-n$, $\nu=\rho+1/2-m$, where m, n are integers, which are different for each term in (34). So we find that three mentioned types of behavior as $\varepsilon\to 0$ are

$$\varepsilon^{\sigma+\rho+d+1-n-m},\qquad \varepsilon^{\sigma-\rho-n+m},\qquad \varepsilon^{\rho-\sigma+n-m}. \tag{38}$$

Note that the powers in the second and the third types are inverse to each other, so if the difference between ρ and σ is not integer we have a singular contribution.

References

[1] J. Maldacena, "The Large N Limit of Superconformal Field Theories and Supergravity", Adv. Theor. Math. Phys. 2 (1998) 231, hep-th/9711200.

[2] S. S. Gubser,I. R. Klebanov and A. M. Polyakov, Gauge Theory Correlators from Non-critical String Theory, Phys. Lett. B428 (1998) 105, hep-th/9802109.

[3] E. Witten, Anti De Sitter Space and Holography, Adv. Theor. Math. Phys. 2 (1998) 253, hep-th/9802150.

[4] I. Ya. Aref'eva and I. V. Volovich, On Large N Conformal Theories, Field Theories in Anti-De Sitter Space and Singletons, hep-th/9803028.

[5] W. Mück and K. S. Viswanathan, Conformal Field Theory Correlators from Classical Scalar Field Theory on AdS_{d+1}, Phys. Rev. D58 (1998) 041901, hep-th/9804035.

[6] D. Z. Freedman, S. D. Mathur, A. Matusis and L. Rastelli, Correlation functions in the CFT_d/AdS_{d+1} correspondence, hep-th/9804058.

[7] I. Aref'eva and I. Volovich, On the Breaking of Conformal Symmetry in the AdS/CFT Correspondence, Phys. Lett. B433 (1998) 49, hep-th/9804182.

[8] W. Mück and K. S. Viswanathan, Conformal Field Theory Correlators from Classical Field Theory on Anti-de Sitter Space II. Vector and Spinor Fields, Phys. Rev. D58 (1998) 106006, hep-th/9805145.

[9] G. Chalmers, H. Nastase, K. Schalm and R. Siebelink, R-Current Correlators in $\mathcal{N}=4$ Super Yang-Mills Theory from Anti-de Sitter Supergravity, hep-th/9805105.

[10] M. Henningson and K. Sfetsos, Spinors and the AdS/CFT correspondence, Phys. Lett. B431 (1998) 63, hep-th/9803251.

[11] H. Liu and A. A. Tseytlin, "$D = 4$ Super Yang-Mills, $D = 5$ gauged supergravity and $D = 4$ conformal supergravity", hep-th/9804083, to appear in NPB.

[12] G. E. Arutyunov and S. A. Frolov, On the origin of supergravity boundary terms in the AdS/CFT correspondence, hep-th/9806216.

[13] W. Mück and K. S. Viswanathan, The Graviton in the AdS-CFT correspondence: Solution via the Dirichlet boundary value problem, hep-th/9810151.

[14] G. E. Arutyunov and S. A. Frolov, Antisymmetric tensor field on AdS_5, hep-th/9807046.

[15] W. S. l'Yi, Generating functionals of correlation functions of p-form currents in AdS/CFT correspondence, hep-th/9809132.

[16] W. S. l'Yi, Correlators of currents corresponding to the massive p-form field in AdS/CFT correspondence, hep-th/9811097.

[17] S. Corley, The massless gravitino and the AdS/CFT correspondence, hep-th /9808184.

[18] A. Volovich, Rarita-Schwinger Field in the AdS/CFT Correspondence, hep-th/9809009.

[19] A. Ghezelbash, K. Kaviani, S. Parvizi and A. Fatollahi, Interacting Spinors - Scalars and ADS/CFT Correspondence, Phys. Lett. B435 (1998) 291, hep-th/9805162.

[20] S. Lee, S. Minwalla, M. Rangamani, N. Seiberg, Three-Point Functions of Chiral Operators in $D = 4$, $\mathcal{N} = 4$ SYM at Large N, hep-th/9806074

[21] P. van Nieuwenhuizen, Supergravity, Phys. Rep. 68C (1981) 189.

[22] H. J. Kim, L. J. Romans and P. van Nieuwenhuizen, Mass spectrum of chiral ten-dimensional $N = 2$ supergravity on S^5, Phys. Rev. D32 (1985) 389.

[23] P. van Nieuwenhuizen, The complete mass spectrum of d=11 supergravity compactified on S^4 and the general mass formula for arbitrary cosets M_4, Class. Quant. Garv. 2 (1985) 1.

[24] S. Ferrara, A. F. Grillo and G. Parisi, The Schadow Operator Formalism for Conformal Algebra. Vacuum Expectation Values and Operator Products, Lett. Nuovo Cim. 4 (1972) 115.

[25] G. E. Arutyunov and S. A. Frolov, Quadratic action for type IIB supergravity on $AdS_5 \times S^5$, hep-th/9811106

[26] G. M. Sotkov and R. P. Zaikov, Conformal invariant two and three-point functions for fields with arbitrary spin, Rept. on Math. Phys. 12 (1977) 375.

[27] A. P. Prudnikov, Yu. A. Brychkov and O. I. Marichev, Integrals and Series, Vol. 3, Nauka, Moscow, 1985